## 'SY Clearstone 2CL' Winter Wheat

## Phil Bruckner and Jim Berg, Winter Wheat Breeding Program, Montana State University Updated 12/2017

SY Clearstone 2CL is a 2-gene CLEARFIELD hard red winter wheat developed by Montana Agricultural Experiment Station in 2012 and licensed exclusively to Syngenta Seeds. SY Clearstone 2CL wheat is very similar to Yellowstone. It is a medium maturing, medium tall, white chaffed wheat with average winter hardiness. It is a high yielding wheat (Tables 1,2) with average test weight and protein (Table 3). SY Clearstone 2CL is resistant to stripe rust and has moderate resistance to stem rust, the latter an improvement over Yellowstone. SY Clearstone 2CL is a medium PPO variety with average mill and above average bake properties (Table 4). <a href="PVP">PVP</a>, Title V has been issued (Certificate# 201300357). Additionally, the CLEARFIELD genes are patented.

Variety	Districts								
	1	2	3	4	5	5	6- Sidney &	All	
	Kalispell	Bozeman	Huntley <sup>2/</sup>	Moccasin <sup>3/</sup>	Conrad <sup>4/</sup>	Havre <sup>5/</sup>	Williston	Locations	
location-years	5	5	30	27	22	15	7	111	
Yellowstone	116.9	92.3	68.4	<u>58.9</u>	<u>78.7</u>	57.6	59.2	69.3	
Northern	<u>120.1</u>	86.9	66.4	58.1	76.4	<u>57.7</u>	53.0	67.7	
SY Clearstone 2CL	117.9	88.7	66.0	58.2	76.4	56.1	51.2	67.3	
CDC Falcon	74.6	71.2	62.8	53.9	71.8	53.8	55.6	61.7	
Decade	50.5	72.0	63.4	55.5	72.6	51.9	51.3	60.8	
LSD (0.05)	19.6	12.7	3.0	2.5	3.1	3.2	ns	2.3	
<b>bold</b> = indicates highest value within a column ns = non-significant									

bold = indicates varieties with values equal to highest variety within a column based on Fisher's Protected LSD (p =0.05)

Table 2. Yield of SY Clearstone 2CL vs. a set of CLEARFIELD varieties: 2015-2017 Intrastate Tests

Variety	Districts							
	1	2	3	4	5	5	6- Sidney &	All  Locations
	Kalispell	Bozeman	Huntley	Moccasin	Conrad	Havre <sup>6/</sup>	Williston	Locations
location-years	2	3	2	3	3	5	4	22
SY Clearstone 2CL	124.6	90.5	116.0	<u>62.0</u>	87.2	63.9	50.6	<u>78.3</u>
WB4623CLP	130.5	90.7	108.8	52.4	77.2	57.9	33.7	71.1
Brawl CL Plus	72.5	71.8	112.3	58.3	86.4	61.6	44.7	68.4
LSD (0.05)	ns	ns	ns	6.4	ns	ns	ns	8.1
		6/ = includes Ft. Benton						

<sup>1/ = 2012-2016</sup> Intrastate and 2013-2017 Off Station tests

<sup>2/</sup> includes data from Forsyth, Fort Smith, Hardin area, Hysham, Lodge Grass, Molt, Rapelje

<sup>3/</sup> includes data from Belt, Denton, Geraldine, Highwood, Winifred

<sup>4/</sup> includes data from Choteau, The Knees, Shelby

<sup>5/</sup> includes data from Ft. Benton, Loma, Turner

Table 3. Agronomic characteristics of SY Clearstone 2CL vs. a set of recommended varieties, 2012-2017<sup>1/</sup>

Variety	Test	Winter	Heading date		Plant	Lodging	Protein	Sawfly	Stripe	Coleoptile
	weight lb/bu	survival %	Julian	Calendar	height in	%	%	cutting %	rust %	length in
location-years	111	5	49		109	20	108	11	8	2
CDC Falcon	59.1	<u>55</u>	160.1	9-Jun	29.7	23	12.6	25	53	2.9
Decade	59.3	50	159.4	8-Jun	31.4	24	12.8	20	71	3.2
Northern	<u>59.5</u>	40	162.5	12-Jun	31.5	23	<u>12.8</u>	15	<u>22</u>	2.5
SY Clearstone 2CL	58.9	33	161.3	10-Jun	33.8	27	12.4	23	34	3.0
Yellowstone	59.4	45	161.4	10-Jun	33.2	23	12.4	21	33	2.7
LSD (0.05)	0.34	12	0.4		0.3	ns	0.2	ns	14	0.2

<sup>1/ = 2012-2016</sup> Intrastate and 2013-2017 Off Station tests

ns = non-significant

**bold** = indicates highest value within a column

bold = indicates varieties with values equal to highest variety within a column based on Fisher's Protected LSD (p =0.05)

Table 4. Mill and bake characteristics of SY Clearstone 2CL vs. a set of varieties, 2012-2016

Variety	PPO 1/	Kernel	Flour				Mixograph	1	Baking		
		hardness	yield	protein	ash	tolerance	mix time	absorption	mix time	absorption	volume
			%	%	%	(1-6)	min	%	min	%	СС
location-years	20	20	20	20	20	20	20	20	20	20	20
Decade Northern	0.285 <b>0.104</b>	75.7 86.2	68.6	11.5	<b>0.42</b> 0.45	<u>4.6</u> 3.5	7.9 4.1	<b>64.9</b> 62.4	19.2 6.2	75.3 72.2	1044 1076
SY Clearstone 2CL	0.300	77.8	<b>69.7</b> 67.8	<u>11.8</u> 11.1	0.43	4.0	6.1	62.4	9.9	72.6	1078
Yellowstone	0.217	78.9	68.9	11.3	0.43	4.4	8.3	64.2	16.0	75.0	1052
LSD (0.05)	0.032	2.4	0.7	0.3	0.01	0.4	0.7	1.2	2.0	1.2	ns

**bold** = indicates highest value within a column

ns = non-significant

1/ low is best for noodles

Phil Bruckner and Jim Berg, Montana State University, Agricultural Experiment Station <a href="http://plantsciences.montana.edu/crops">http://plantsciences.montana.edu/crops</a>

**bold** = indicates varieties with values equal to highest variety within a column based on Fisher's Protected LSD (p =0.05)